

## Use of lipid-lowering medicinal herbs during pregnancy: A systematic review on safety and dosage

Hojjat Rouhi-Boroujeni<sup>(1)</sup>, Esfandiar Heidarian<sup>(2)</sup>, Hamid Rouhi-Boroujeni<sup>(3)</sup>, Minasadat Khoddami<sup>(4)</sup>, Mojgan Gharipour<sup>(5)</sup>, Mahmoud Rafeian-Kopaei<sup>(6)</sup>

### Review Article

#### Abstract

**BACKGROUND:** Hyperlipidemia is one of the important diseases in pregnancy that causes fetal abnormalities during pregnancy and after the birth. Unfortunately, the usual anti-fat drugs are associated with high morbidity in fetus and due to people's inclination towards taking herbs, it is required to identify side effects of medicinal herbs in pregnancy. The aim of this study was to present hypolipidemic herbs that would not any complications for mother and fetus.

**METHODS:** In this review article, the major electronic databases such as EBSCO, Central Register of Controlled Trials (CENTRAL), China Network Knowledge Infrastructure (CNKI), Cochrane, Google scholar, MEDLINE, SciVerse, Scopus, and Web of Science were searched using the key words “herbal” and “hyperlipidemia”, “herbal” and “pregnancy” matched by MeSH from their respective inception till September, 2016. Total of 1723 publications (145 review articles, 855 original research articles, and 723 abstracts) about the effect of herbals on hyperlipidemia and 682 publications (200 abstracts, 423 original research articles, and 59 review articles) about the effect of herbals in pregnancy were retrieved. At the end, a list of medicinal plants effective on hyperlipidemia alongside their effects on pregnancy was developed. Finally, the plants effective on hyperlipidemia and safe during pregnancy were determined and their dosage, complications, mechanism of action, and side effects were reported.

**RESULTS:** A total of 110 effective herbs on hyperlipidemia were identified and complications of 95 plants in pregnancy were studied. At last, among the 55 selected plants effective on hyperlipidemia and examined for pregnancy, we reported 12 herbs with their dosage and special considerations that can be used to treat hyperlipidemia during pregnancy.

**CONCLUSION:** Some medicinal plants can be used to treat hyperlipidemia during pregnancy without any significant side effects both on mother or fetus.

**Keywords:** Hyperlipidemias, Pregnancy Outcome, Fertility, Dyslipidemia, Herbals, Medicinal Plants, Oxidative Stress

*Date of submission:* 15 Jan. 2017, *Date of acceptance:* 21 Mar. 2017

#### Introduction

Exposure to elevated levels of cholesterol and oxidative stress due to products of cholesterol metabolism during fetal period has been shown to result in programmed death of fetal arterial cells with a predisposition to atherosclerosis later in life.<sup>1</sup> Commonly, during reproductive years (about 2 decades), risk of cardiovascular diseases reduces.

Besides, lipid and lipoproteins is not been measured routinely during pregnancy as gestational dyslipidemia is considered physiologic with little clinical significance.<sup>2</sup> However, recent discoveries of fatty streaks in the aorta of 6-month-old fetuses and also evidences of aortic atherosclerosis in autopsy of deceased infants with normal levels of cholesterol born to mothers with

1- PhD Candidate, Student Research Committee, Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

2- Professor, Clinical Biochemistry Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

3- Pulmonologist, Clinical Biochemistry Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

4- Pharmacist, Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

5- PhD Candidate, Cardiac Rehabilitation Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

6- Professor, Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

Correspondence to: Mahmoud Rafeian-Kopaei, Email: rafeian@yahoo.com

hypercholesterolemia, has highlighted the importance of correcting or preventing maternal dyslipidemia for the benefit of the mother and the child.<sup>3</sup> Currently, no reference standards exist for lipid parameters during pregnancy, although it is well-known that pregnancy is a state of insulin resistance, which is reflected by lipoprotein lipid profiles. Pregnancy-related hypertriglyceridemia is rare, but it can be life threatening in some patients with genetic susceptibility. Complications can include acute pancreatitis, hyperviscosity syndrome, and potentially preeclampsia. Overweight and obese women are significantly more likely to exceed the pregnancy-related weight gain recommendations. Women with gestational diabetes and/or preeclampsia are also at increased risk for elevated triglyceride levels, development of chronic hypertension, recurrent gestational diabetes and/or overt diabetes, recurrent preeclampsia, and development of albuminuria later in life.<sup>4</sup>

Two registered clinical trials are currently evaluating the effects of lipophilic statins to prevent preeclampsia in pregnancy. The true risk of congenital anomalies caused by statins in pregnancy has not been well confirmed in humans yet. However, because statins are category X, they should only be used in a research setting during pregnancy until more information is available. Fenofibrate has been assigned to pregnancy category C by the Food and Drug Administration (FDA). Fenofibrate should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. The side effects of statins and other antihyperlipidemic drugs in animal models of pregnancy showed delayed delivery, increased postimplantation loss, decreased litter size and pup birth weight, 40% pup survival rate, 4% neonate survival, no pup survival to weaning, and increased incidence of spina bifida, abortion, and fetal skeletal abnormalities (domed head, hunched shoulders, rounded body, abnormal chest, kyphosis, stunted fetuses, altered skeletal formation of ribs, sternbrae, vertebrae, and palatine). Delayed delivery, decreased live births, and death of 17% of fetuses occurred at doses 18 times higher than the maximum human dosage. In addition, studies on animal reproductive system with doses 7 to 10 times higher than the recommended human dosage based on body surface area (BSA) have demonstrated to have embryocidal and teratogenic effects.<sup>5,6</sup>

Lifestyle changes and glycemic control should be instituted if necessary. During pregnancy, a bile acid sequestrant can safely treat elevated cholesterol


levels. Women must be educated about dietetic measures and body mass reduction even in preconception period. In addition, during pregnancy, mothers must be monitored and due to risk of pancreatitis in case of triglyceride above 11.5 mmol/l, other therapy options must be taken into account. In the last trimester of pregnancy, severe hypertriglyceridemia associated with pancreatitis can be treated with omega-3 fatty acids, parenteral nutrition, plasmapheresis, and other lipid-lowering agents.<sup>7</sup>

The use of herbal medicines has been increasing in many developing and industrialized countries. More and more pregnant women are using herbal remedies to treat pregnancy-related problems due to cost-effectiveness of therapy and easy access to these products.<sup>8</sup>

To date, over 200 plants have been recommended for treatment of hyperlipidemia. As with chemical drugs, medicinal plants can cause permanent damage to fetus. Therefore, despite people's willingness to use medicinal plants, certain precautions with these plants should be taken into account. In addition, couples are likely to use these plants on the verge of fertility to treat hyperlipidemia or other disorders.<sup>9,10</sup> Therefore, it is highly necessary for both physicians and patients to know which plants have optimal effects on hyperlipidemia during and before pregnancy without having side effects.<sup>11</sup> The aim of this review article was to investigate the effect of plants on hyperlipidemia and plant-based side effects in pregnancy and fertility as well as to introduce the plants that are effective on hyperlipidemia during pregnancy.

## Materials and Methods

In this study, 2405 publications (204 review articles, 1278 original full text articles, and 923 abstracts) were retrieved. The major electronic databases including Web of Science, Scopus, PubMed, Google scholar, MEDLINE, EBSCO, China Network Knowledge Infrastructure (CNKI), and Cochrane Central Register of Controlled Trials (CENTRAL) were searched from their respective inceptions till September 2016. To identify herbs used to treat hyperlipidemia the following keywords were used and matched by the MeSH: “herbal in hyperlipidemia”, “botany in hyperlipidemia”, “herbal therapy in hypertriglyceridemia”, “systematic review of herbal in hypercholesterolemia”, “herbal medicine for hypercholesterolemia”, “herbal with anti-lipid effect”, “natural remedies for hyperlipidemia”, “herbal therapy for atherosclerosis and “hypolipidemic diet”.



**Figure 1.** Searching and data extraction was based on the Cochrane protocol and checklist for review

Total of 1723 publications (145 review articles, 855 original research articles, and 723 abstracts) were analyzed and their findings are registered in checklist 1.

We selected herbal drugs based on safety in pregnancy. All steps for searching and data extraction was based on the Cochrane protocol and checklist for systematic review (Figure 1).

In addition, to find evidence on the efficacy of herbals in pregnancy, fertility and infertility, 692 publications (200 abstracts, 423 original research articles, and 59 review articles) were analyzed. The headings that were used included “herbal in pregnancy”, “phytomedicine in pregnancy”, “side effects of herbal in pregnancy”, “herbals in pregnancy and lactation”, “herbal therapy in fertility”, “herbal therapy in infertility”, “herbal in fertility”, “herbal in infertility”, “phytomedicine in infertility”, “botany in pregnancy”, “medicinal plants in fertility”, “Chinese herbal in pregnancy”, “review of herbal in pregnancy”, and “Ayurvedic herbal in pregnancy”. The results of this investigation were registered in checklist 2. A plant was included in the analysis if its name appeared in at least two publications. Then, the plants effective on hyperlipidemia, fertility, and pregnancy were determined after the two checklists were integrated (Table 1). Finally, the plants effectiveness on hyperlipidemia and safety during pregnancy were determined and after analysis of 110 publications,

their dosage, complications, mechanisms of action, and side effects were reported (Table 2).

## Results

A total of 110 plants have been reported to be effective on hyperlipidemia and 95 plants were reported to be effective on fertility and pregnancy. Overall, 12 and 55 plants have been reported to be effective on lipid and safe during pregnancy, respectively. The potential side effects, dosage, and special considerations regarding these plants are shown in table 2. Moreover, 21 plants could be used in normal diet during pregnancy but were not recommended as medicinal plants.

## Discussion

Hyperlipidemia can affect maternal and fetal health. Many side effects of chemical drugs on mother and fetus have led to prevention of their use during pregnancy. In this study, we found that the effective medicinal plants on hyperlipidemia contributed greatly to reducing oxidative stress via their antioxidant properties in addition to directly exerting hypolipidemic effects.

Reactive oxygen species cause damage to the structure of different cells and tissues including heart and vessels. Napoli *et al.* demonstrated that low levels of superoxide dismutase (SOD) in pregnant rabbits that had hyperlipidemia for over six months led to formation of fatty streaks in the aortic arch in their fetus.<sup>11</sup>

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
<i>Achillea millefolium</i> <sup>12-14</sup>	Yarrow	Asteraceae	Leaf	May interfere with spermatogenesis	-	-	Abortifacient, emmenagogue	Reduces fetal weight, increases placental weight, neurotoxic component, Potential harmful.	Prohibited in pregnancy, even with nutritional values
<i>Allium cepa</i> <sup>15,16</sup>	Onion	Liliaceae	Leaf, bulb	-	-	-	-	-	Lower risk of spontaneous preterm delivery
<i>Allium sativum</i> <sup>17-20</sup>	Garlic	Liliaceae	Leaf, bulb	--	Minimal risk – third trimester, crosses into the amniotic fluid	-	-	Potential abortifacient, emmenagogue, uterine stimulant	In clinical and animal studies, at doses lower than 1 g, no complications were seen This plant was used to lower preeclampsia and hyperlipidemia during pregnancy Lower risk of spontaneous preterm delivery
<i>Aloe vera</i> <sup>21-25</sup>	Cap aloe	Liliaceae	Leaf	Antifertility effect in male	-	Potentially nephrotoxic, potential hepatic dysfunction	Potentially genotoxic, mutagenic, carcinogenic	Potential abortifacient, emmenagogue Aloe vera gel – minimal risk	Prohibited in pregnancy, even with nutritional values
<i>Anethum graveolens</i> <sup>26-30</sup>	Dill	Apiaceae	Leaf, seed	Induces infertility without any effect on oocyte structure, decreases sexual potency and spermatogenesis in males	Uterine muscles of rat contracted in the presence of dill	-	-	-	Induction of labor
<i>Apium graveolens</i> <sup>31</sup>	Celery	Umbellifera	Leaf	-	Uterine stimulant, abortifacient and emmenagogue	-	-	-	-

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
<i>Artemisia vulgaris</i> <sup>32,33</sup>	Mugwort	Compositae	Leaf	-	Emmenagogue and abortifacient effects	-	-	-	Prohibited in pregnancy, even with nutritional values
<i>Arctium loppa</i> <sup>8</sup>	Burdock	Compositae	Root	-	Oxytotic and uterine stimulant action	-	-	-	Prohibited in pregnancy, even with nutritional values
<i>Avena sativa</i> <sup>34-36</sup>	Oats	Gramineae	Fruit	-	-	-	-	-	No data available
<i>Berberis vulgaris</i> <sup>37</sup>	Barberry	Berberidaceae	Root and fruit	-	-	-	May cause newborn jaundice (kernicterus)	Uterine stimulant	-
<i>Boswellia carterii</i> <sup>38-40</sup>	Indian tree	Burseraceae	Resin	An aphrodisiac and a fertility promoting agent, increases sperm motility and sperm density	-	-	-	-	There is lack of evidence on safe use of boswellia during pregnancy and lactation
<i>Calendula officinalis</i> <sup>4,5,25,41</sup>	Marigold-calendula	Compositae	Flower	Spermicide, anti-blastocyst	-	-	Uterotonic effect	Emmenagogue, potential abortifacient, estrogenic	Topical-unknown
<i>Chicorium intybus</i> <sup>42</sup>	Chicory	Compositae	Root	-	Reduces body weight, weight gain, body length and serum free fatty acids, uterine contractions	-	-	-	Prohibited in pregnancy, even with nutritional values
<i>Citrus limon</i> <sup>43-45</sup>	Lemon	Rutaceae	Fruit	Anti-fertility effect in men	-	-	-	-	Lemon inhalation can be effective in reducing nausea and vomiting of pregnancy

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
Cinnamomum verum <sup>46-53</sup>	Cinnamon	Lauraceae	Bark	Significant increase in reproductive organ weights, sperm motility, sperm count	-	-	-	Emmenagogue effects	Unsafe for therapeutic use during pregnancy It is not recommended to be used in food during pregnancy A uterine stimulant in high doses, but quite safe as a culinary herb; avoid the essential oil completely
Citrus paradise <sup>54,55</sup>	Grapefruit	Rutaceae	Fruit	-	Safe	-	-	-	At edible amounts during pregnancy, it is used as an effective antioxidant and fibrous food, over once daily is not recommended and interactions with other drugs and supplements should be taken into account
Coffea Arabica <sup>56-63</sup>	Arabica coffee	Rubiaceae	Seed	-	Spontaneous abortion, increased risk of stillbirth, low birth weight infants	-	Teratogenic compounds, impairs trace mineral absorption in fetus	Harmful to the fetus (crosses the placenta)	Three cups of coffee throughout the day possibly safe
Commiphora mukul <sup>16,25,64</sup>	Guggul	Burseraceae	Gum	-	-	-	-	Potential abortifacient, Emmenagogue, uterine stimulant	Prohibited in pregnancy, even with nutritional values
Cornus mas <sup>65-67</sup>	Cran berry	Cornaceae	Fruit	-	-	-	-	-	Herbal compendium reported that cranberry is of minimal risk when consumed safe in food quantities It is used to treat uterine tract infections during pregnancy

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
<i>Crataegus microphylla</i> C. Koch <sup>68-69</sup>	Howthorn	Rosaceae	Leaf, fruit	-	-	-	-	Uterine activity	-
<i>Dioscorea nipponica</i> <sup>70</sup>	Wild yam	Dioscoreaceae	Rhizome	-	Contractile agonist for the uterus, abortion	-	-	-	-
<i>Eleutherococcus</i> <sup>71-74</sup>	Ginseng	Araliaceae	Rhizome	-	-	-	-	-	Panax ginseng should be consumed with caution during pregnancy, especially during the first trimester
<i>Equisetum arvense</i> <sup>74</sup>	Horsetail	Equisetaceae	-	-	-	-	-	May cause autism	There are few studies about this plant and it is better not to be used in pregnancy
<i>Eucalyptus globulus</i> <sup>75,76</sup>	Eucalyptus	Myrtaceae	Leaf	Decreases fertility in male	-	-	-	-	There has been no adverse outcome in mice injected on days 6 and 15 of gestation There has been no evidence of adverse reproductive effects of eucalyptus oil in humans
<i>Ficus carica</i> <sup>77</sup>	Fig	Moraceae	Leaf and fruit	-	-	-	-	-	Topically, it is safe Fresh or dried fig fruit is likely safe in amounts found in food, but there is not enough information to know if it is safe in the larger amounts that are used as medicine Lower risk of spontaneous preterm delivery

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
Ginco biloba <sup>78-80</sup>	Ginkgo	Ginkgoaceae	Leaf	-	malformations including round shaped eye and orbits, syndactyly, malformed pinnae, nostrils, lips and jaws.	Unsafe when adulterated with colchicine, antiplatelet, emmenagogue, hormonal changes	Ginkgo leaf has antiplatelet activity, which may be of concern during labor as ginkgo use could prolong bleeding time	Emmenagogue, hormonal changes	Prohibited in pregnancy, even with nutritional values
Glycine soja <sup>81,82</sup>	Soy	Leguminosae	Seed	-	-	-	-	-	Prohibited in pregnancy, even with nutritional values
Glycyrrhiza glabra <sup>83,84</sup>	Licorice	Leguminosae	Root	-	-	Likely to be born before 38 weeks of gestation, risk of pre-term pregnancy (before 37 weeks), does not affect birth weight, does not affect maternal blood pressure	-	Potential abortifacient, emmenagogue, uterine stimulant, causes high prolactin and estrogen levels, risk of pre-term pregnancy (before 37 weeks), does not affect birth weight	-
Hibiscus sabdariffa <sup>85,86</sup>	Hibiscus	Malvaceae	Flower	-	-	-	-	Decrease both pregnancy weight gain and postpartum weight loss, decrease maternal fluid and food intake with increased plasma sodium and corticosterone concentration	There is some evidence that hibiscus might start menstruation, and this could cause a miscarriage Aromatic ketones may present some hazard



**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
<i>Lavandula stoechas</i> <sup>87-89</sup>	Lavender	Labiatae	Leaf	-	-	-	-	Emmenagogue effects	Lavender oil had estrogenic and anti-androgenic activities Due to its purported properties as an emmenagogue, excessive internal use should be avoided during pregnancy; however, there is no definitive evidence in this area Safe in pregnancy
<i>Malus orientalis</i> <sup>9</sup>	Apple	Rosaceae	Fruit	-	-	-	--		
<i>Medicago sativa</i> <sup>25,68,90,91</sup>	Alfalfa	Leguminaceae	Leaf	Antifertility in man	Estrogenic activity			Emmenagogue, anti-gonadotrophic activity	Minimal risk in food
<i>Nigella sativa</i> <sup>92,93</sup>	Black cumin	Ranunculaceae	Seed	<i>Nigella sativa</i> oil L. (Ranunculaceae) and Cinnamon zeylanicum J. Presl (Lauraceae) were found to enhance fertility	Stimulation of uterine contractions, abortion	-	-	-	-
<i>Oenothera bienni</i> <sup>94-96</sup>	Evening primrose	Onagraceae	Seed		Teratogenic and induces labor <sup>15</sup>	May induce labor but effectiveness is unclear, increased risk of pregnancy complication (evidence level 1b), prolonged rupture of membranes, oxytocin augmentation, arrest of descent, vacuum extraction			Oral administration of evening primrose oil from the 37 <sup>th</sup> gestational week until birth does not shorten gestation or decrease the overall length of labor Further, the use of orally administered evening primrose oil may be associated with an increase in the incidence of prolonged rupture of membranes, oxytocin augmentation, arrest of descent, and vacuum extraction

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
Ocimum basilicom <sup>97</sup>	Basil	Labiatae	Leaf	-	-	-	-	Emmenagogue, abortifacient, mutagenic	-
Peganum harmala <sup>98</sup>	Harmala	Zygophyllaceae	Seed	-	-	-	-	-	Prohibited in pregnancy, even with nutritional values
Persea Americana <sup>99</sup>	Avocado	Lauraceae	Seed, fruit	-	-	-	-	-	There is not enough reliable information about the safety of taking avocado as medicine if you are pregnant or breast-feeding, stay on the safe side and stick to food amounts
Petroselinum crispum <sup>100</sup>	Parsley	Umbelliferae	Leaf	-	Abortifacient	-	-	Emmenagogue, estrogenic, uterine stimulant constituent	-
Plantago psyllium <sup>101-104</sup>	Plantain	Plantaginaceae	Leaf, seed	-	-	-	-	-	Psyllium powder could significantly decrease the number of surgeries resulting from anorectal complications, hemorrhoid diseases, anal fissure and constipation It is in concordance of several other studies which emphasized the effect of fiber in diet on preventing constipation in the course of pregnancy If used in low amounts in diet, it causes no problem
Purtolaca oleraceae <sup>105</sup>	Purslane	Purtulaceae	Leaf	Antifertility effect in male rat	-	Abortifacient	-	-	Sweet cherry is safe for pregnant and breast-feeding women in food amounts, but larger medicinal amounts should be avoided until more is known
Pronus avium <sup>106</sup>	Cherry	Rosaceae	Fruit, cherry tails	-	-	-	-	-	

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
<i>Punica granatum</i> <sup>107</sup>	Pomegranate	Punicaceae	Fruit, leaf	-	-	-	-	-	Use cautiously in pregnant and breastfeeding women, due to a lack of safety data Although some animal studies show that pomegranate may induce abortion, consuming pomegranate as a food is likely safe during pregnancy There is little information available on the topical use (application to the skin) of pomegranate during pregnancy and breastfeeding
<i>Rhus coriaria</i> L. <sup>108</sup>	Sumac	Anacardiaceae	Fruit	-	-	-	-	-	Cautionary herb during pregnancy
<i>Solanum lycopersicum</i> <sup>109</sup>	Tomato	Solanaceae	Fruit	-	-	-	-	-	Safe in pregnancy
Tea <i>sinensis</i> <sup>60,67,110-114</sup>	Tea, green tea	Theaceae	Leaf	-	Spontaneous abortion, increased risk of stillbirth, low birth weight infants	-	-	Harmful to the fetus	Three cups or more of tea per day was associated with an increased risk of spina bifida
<i>Taraxacum officinale</i> <sup>68,115</sup>	Dandelion	Compositae	Root, leaf	-	-	-	-	-	Minimal risk in food amounts No negative effects on humans have been reported during pregnancy or lactation, in children, or in combination with pharmaceutical drugs

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
<i>Terminalia chebul</i> <sup>116</sup>	Haritaki	Combretaceae	Fruit	-	-	-	-	-	There is some evidence that <i>Terminalia arjuna</i> is possibly unsafe during pregnancy The safety of the other two species during pregnancy is unknown. It is best to avoid using any terminalia species Topically, it is safe
<i>Thymus vulgaris</i> <sup>117-119</sup>	Thyme	Labiataeae	Leaf	Decreases fertility in male	-	-	-	Emmenagogue, abortifacient	
<i>Trigonella foenum</i> <sup>25,120,121</sup>	Fenugreek	Leguminosae	Seed	-	-	Pseudo-maple syrup urine disease	Potential abortifacient Uterine stimulant	Emmenagogue	Minimal risk in food
<i>Urtica dioica</i> <sup>122-124</sup>	Nettle	Urticaceae	Root, leaf	Increasing fertility in women and men, increase the quality of spermatozoa and inhibits nicotine-induced adverse effects on sperm parameters.	Induce uterine stimulation	-	-	-	Use of nettle should be avoided during pregnancy or lactation
<i>Vitex doniana</i> <sup>125-128</sup>	Black plum	Lamiaceae	Fruit	Due to treatment of hyperprolactinemia, premenstrual syndrome, abnormal menstrual cycle, amenorrhea, mastodynia, this herb can induce fertility in woman	Uterine muscle contractions and also potentiated the contractile effects of prostaglandins, ergometrine and oxytocin	-	-	-	Use of vitex agnus cactus (VAC) should be avoided during pregnancy or lactation

**Table 1.** Study of hypolipidemic plants and their effects on fertility and pregnancy (continue)

Scientific name	Common name	Family	Part of use	Pre pregnancy effects	Strong scientific evidence	Good scientific evidence	Fair scientific evidence	Weak scientific evidence	End result or explain certain points
<i>Vitis vinifera</i> <sup>129,130</sup>	Grape	Vitaceae	Fruit, leaf, seed	-	-	-	-	-	Typically, it is safe The grape seed extract was non-mutagenic in mice There are no adverse outcomes in mice
<i>Withania somnifera</i> (L.) Dunal <sup>131,132</sup>	Winter cherry -	Solanaceae	Fruit	Increasing sperm motility and treatment of libido, sexual performance, sexual vigor, and penile erectile dysfunction	Abortion	-	-	-	Prohibited in pregnancy, even with nutritional values
<i>Zingiber officinalis</i> <sup>133-141</sup>	Ginger	Zingiberaceae	Root	-	Minimal risk (up to 1000 mg of dried ginger per day), unlikely cause of spontaneous abortion	Does not increase rates of major malformations	Non-mutagenic, non-teratogenic Mutagenic constituents Anti-mutagenic constituents Potential embryotoxicity	Non-teratogenic.	Ginger could be considered a harmless and possibly effective alternative option for women suffering from nausea and vomiting of pregnancy (NVP)
<i>Zizyphus vulgaris</i> <sup>142,143</sup>	Jujuba	Rhamnaceae	Fruit	Antifertility/contraception, antisteroidogenic activity and hence fertility in adult female mice It was found to arrest the normal estrus cycle of adult female mice at diestrus stage and reduced the wet weight of ovaries significantly Hematological profiles, biochemical estimations of whole blood and serum remained unaltered in extract-treated mice	Consumer safety in pregnancy has not been established	-	-	-	-

**Table 2.** Hypolipidemic herbs that seem safe in pregnancy

Common name	Dosage	Side effects	Special notification
Onion <sup>144</sup>	50 g of fresh onions or 5 g of dried drug	No health hazards or side effects are known in conjunction with the proper administration of designated therapeutic dosages The intake of large quantities can lead to stomach complaints	Popular: pressed juice and onion syrup, made of 500 g onions, 500 g water, 100 g honey and 350 g sugar
Garlic <sup>145-147</sup>	300 mg dry powder or 2 g fresh garlic	Abdominal discomfort, nausea, vomiting, diarrhea and a feeling of fullness have occurred with garlic therapy	Fresh garlic is not recommended in pregnancy
Lemon <sup>147,148</sup>	1 g dry powder infuse	No health hazards or side effects are known in conjunction with the proper administration of designated therapeutic dosages	Avoid the use of commercial liquid products because additional ingredients or fake lemon
Cranberry <sup>147,149</sup>	10 ripe fruit twice a day after meal, 10 ml cranberry juice twice daily after meal	Mild stomach upset and diarrhea	
Fig <sup>147,150</sup>	5 fruit twice daily	No health hazards or side effects are known in conjunction with the proper administration of designated therapeutic dosages	It is better to be soaked in water
Apple <sup>147,151</sup>	3 fruit/day	No health hazards or side effects are known in conjunction with the proper administration of designated therapeutic dosages	Apple seeds are highly toxic, avoid taking it
Psyllium <sup>147,152</sup>	1g in 100 ml water twice daily	Allergic reactions ranging from sneezing to chest congestion and wheezing were reported in three nurses after psyllium use	The dose should be taken 30 min to one hour after taking other medications
Cherry <sup>147,153</sup>	2-5 g dry powder, 10-15 fresh fruit	No health hazards or side effects are known in conjunction with the proper administration of designated therapeutic dosages	Storage: pomegranate should be sealed in containers and protected from moisture
Pomegranate <sup>147,154</sup>	10 ml of juice twice a day or 20 g pomegranate seeds twice a day or 1 tablet/day (90 mg ellagic acid)	No health hazards or side effects are known in conjunction with the proper administration of designated therapeutic dosages	-
Tomato <sup>147,155</sup>	Three tomatoes a day, or 1 g dry powder three times/day	No health hazards or side effects are known in conjunction with the proper administration of designated therapeutic dosages	
Grape <sup>147, 156,157</sup>	10 g fresh fruit, 1 g dry powder	No health hazards or side effects are known in conjunction with the proper administration of designated therapeutic dosages	
Ginger <sup>132, 147,158</sup>	1 g dry powder/day	Increases appetite	Not recommended more than 1 g/day

Besides, Rumbold et al.<sup>159</sup> and Mistry et al.<sup>160</sup> investigated the role of antioxidants in reducing oxidation of fatty acids and decrease in fatty streaks in fetal heart. Clinical trials have demonstrated that oxidative stress due to hyperlipidemia during pregnancy causes circulatory disorders in fetus, delayed fetal development, and increased eclampsia.

Moreover, Jenkins et al. reported that there was significant association between decrease in SOD and increase in miscarriage in pregnant women with hyperlipidemia.<sup>161</sup> According to the evidence, the antioxidant properties of the plants are due to polyphenols, flavonoids, flavonols, gallic acid, and anthocyanins that cause decrease in malondialdehyde (MDA) and increase in SOD, catalase, and glutathione peroxidase (GPX).<sup>162</sup>

Some of the potent antioxidants that not only improve hyperlipidemia in pregnant women but also play a role in protecting the cardiovascular system of the fetus and the mother are as follows: allyl propyl disulphide, sterol, saponin, and quercetin in onion, allicin, allyl di- and trisulphide, alliin, ajones, vinylthiins in garlic, cyanidin, malvidin, peonidin, pelargonidin, petunidin and bioflavonoids in cranberry, bioflavonoids, polyphenols and triterpenoids, quercetin, catechin, phloridzin and chlorogenic acid in apple, anthocyanin (cyanidin-3-rutinoside) and phenolic compounds (flavonol p-coumaroylquinic acid) in cherry, punicalagins, ellagic acid, unicic acid, phytoestrogens, and anthocyanins in pomegranate, vitamins A, B, and E and lycopene in tomato, anthocyanin, vitamins A and E, polyphenols, oligostibenes, and ampelopsins in red grapes and zingiberene, curcumen, bisabolene, gingerols, and zerumbone in ginger.<sup>163</sup>

### Conclusion

There are effective plants that can play a fundamental role in cardiovascular health in mother and fetus by reducing hyperlipidemia.

### Acknowledgments

This article has been derived from the PhD thesis of the first author and financially supported by the Research Deputy of Shahrekord University of Medical Sciences, Shahrekord, Iran.

### Conflict of Interests

Authors have no conflict of interests.

### References

- Dukic A, Zivancevic-Simonovic S, Varjadic M, Dukic S. Hyperlipidemia and pregnancy. *Med Pregl* 2009; 62(Suppl 3): 80-4.
- Laelago T, Yohannes T, Lemango F. Prevalence of herbal medicine use and associated factors among pregnant women attending antenatal care at public health facilities in Hossana Town, Southern Ethiopia: Facility based cross sectional study. *Arch Public Health* 2016; 74: 7.
- Potter JM, Nestel PJ. The hyperlipidemia of pregnancy in normal and complicated pregnancies. *Am J Obstet Gynecol* 1979; 133(2): 165-70.
- Maymunah AO, Kehinde O, Abidoye G, Oluwatosin A. Hypercholesterolaemia in pregnancy as a predictor of adverse pregnancy outcome. *Afr Health Sci* 2014; 14(4): 967-73.
- Kusters DM, Hassani LH, van de Post JA, Wiegman A, Wijburg FA, Kastelein JJ, et al. Statin use during pregnancy: A systematic review and meta-analysis. *Expert Rev Cardiovasc Ther* 2012; 10(3): 363-78.
- Godfrey LM, Erramouspe J, Cleveland KW. Teratogenic risk of statins in pregnancy. *Ann Pharmacother* 2012; 46(10): 1419-24.
- Ali-Shtayeh MS, Jamous RM, Jamous RM. Plants used during pregnancy, childbirth, postpartum and infant healthcare in Palestine. *Complement Ther Clin Pract* 2015; 21(2): 84-93.
- John LJ, Shantakumari N. Herbal medicines use during pregnancy: A review from the Middle East. *Oman Med J* 2015; 30(4): 229-36.
- Rouhi-Boroujeni H, Rouhi-Boroujeni H, Gharipour M, Mohammadzadeh F, Ahmadi S, Rafieian-Kopaei M. Systematic review on safety and drug interaction of herbal therapy in hyperlipidemia: A guide for internist. *Acta Biomed* 2015; 86(2): 130-6.
- Rouhi-Boroujeni H, Rouhi-Boroujeni H, Heidarian E, Mohammadzadeh F, Rafieian-Kopaei M. Herbs with anti-lipid effects and their interactions with statins as a chemical anti- hyperlipidemia group drugs: A systematic review. *ARYA Atheroscler* 2015; 11(4): 244-51.
- Napoli C, Witztum JL, Calara F, de Nigris F, Palinski W. Maternal hypercholesterolemia enhances atherogenesis in normocholesterolemic rabbits, which is inhibited by antioxidant or lipid-lowering intervention during pregnancy: an experimental model of atherogenic mechanisms in human fetuses. *Circ Res* 2000; 87(10): 946-52.
- Montanari T, de Carvalho JE, Dolder H. Antispermatic effect of *Achillea millefolium* L. in mice. *Contraception* 1998; 58(5): 309-13.
- Millet Y, Jouglard J, Steinmetz MD, Tognetti P, Joanny P, Arditti J. Toxicity of some essential plant oils. Clinical and experimental study. *Clin Toxicol* 1981; 18(12): 1485-98.
- Bonkovsky HL, Cable EE, Cable JW, Donohue SE, White EC, Greene YJ, et al. Porphrogenic properties of the terpenes camphor, pinene, and thujone (with a note on historic implications for

- absinthe and the illness of Vincent van Gogh). *Biochem Pharmacol* 1992; 43(11): 2359-68.
15. Ige SF, Akhigbe RE. Common onion (*Allium cepa*) extract reverses cadmium-induced organ toxicity and dyslipidaemia via redox alteration in rats. *Pathophysiology* 2013; 20(4): 269-74.
  16. McGuffin M, Hobbs C, Upton R, Goldberg A. *Botanical safety handbook*. Boca Raton, FL: CRC Press; 1997.
  17. Bercaw J, Maheshwari B, Sangi-Haghpeykar H. The use during pregnancy of prescription, over-the-counter, and alternative medications among Hispanic women. *Birth* 2010; 37(3): 211-8.
  18. El-Sayyad HI, Abou-El-Naga AM, Gadallah AA, Bakr IH. Protective effects of *Allium sativum* against defects of hypercholesterolemia on pregnant rats and their offspring. *Int J Clin Exp Med* 2010; 3(2): 152-63.
  19. Ziaei S, Hantoshzadeh S, Rezasoltani P, Lamyian M. The effect of garlic tablet on plasma lipids and platelet aggregation in nulliparous pregnant women at high risk of preeclampsia. *Eur J Obstet Gynecol Reprod Biol* 2001; 99(2): 201-6.
  20. Mennella JA, Johnson A, Beauchamp GK. Garlic ingestion by pregnant women alters the odor of amniotic fluid. *Chem Senses* 1995; 20(2): 207-9.
  21. Oyewopo A, Oremosu A, Akang E, Noronha C, Okanlawon A. Effects of Aloe Vera (*Aloe barbadensis*) aqueous leaf extract on testicular weight, sperm count and motility of adult male Sprague-Dawley rats. *Journal of American Science* 2011; 7(4): 31-4.
  22. Seetharam YN, Sujeeth H, Jyothishwaran G, Barad A, Sharanabasappa G, Umareddy B, et al. Antifertility effect of ethanolic extract of Amalakyadi churna in male albino mice. *Asian J Androl* 2003; 5(3): 247-50.
  23. Suzuki I, Saito H, Inoue S, Migita S, Takahashi T. Purification and characterization of two lectins from *Aloe arborescens* Mill. *J Biochem* 1979; 85(1): 163-71.
  24. Nath D, Sethi N, Singh RK, Jain AK. Commonly used Indian abortifacient plants with special reference to their teratologic effects in rats. *J Ethnopharmacol* 1992; 36(2): 147-54.
  25. Farnsworth NR, Bingel AS, Cordell GA, Crane FA, Fong HH. Potential value of plants as sources of new antifertility agents I. *J Pharm Sci* 1975; 64(4): 535-98.
  26. Malihezaman M, Mojaba M, Elham H, Farnaz G, Ramin M. Anti-fertility effects of different fractions of *Anethum graveolens* L. extracts on female rats. *Afr J Tradit Complement Altern Med* 2012; 9(3): 336-41.
  27. Monsefi M, Ghasemi M, Bahaoddini A. The effects of *Anethum graveolens* L. on female reproductive system. *Phytother Res* 2006; 20(10): 865-8.
  28. Monsefi M, Ghasemi M, Bahaoddini A. The effects of *Anethum graveolens* L. on female reproductive system of rats. *DARU J Pharm Sci* 2006; 14(3): 131-5.
  29. Monsefi M, Ghasemi A, Alae S, Aliabadi E. Effects of *Anethum graveolens* L. (dill) on oocyte and fertility of adult female rats. *J Reprod Infertil* 2015; 16(1): 10-7.
  30. Akbari M, Javadnoori M, Siahpoosh A, Afshari P, Haghghi MH, Lake E. Comparison the effect of *Anethum graveolens* and oxytocin on induction of labor in term pregnancy: A randomized clinical trial. *Jundishapur J Nat Pharm Prod* 2016; 11(1): e27876.
  31. Wilkinson JM. What do we know about herbal morning sickness treatments? A literature survey. *Midwifery* 2000; 16(3): 224-8.
  32. Tigno XT, Gumila E. In vivo microvascular actions of *Artemisia vulgaris* L. in a model of ischemia-reperfusion injury in the rat intestinal mesentery. *Clin Hemorheol Microcirc* 2000; 23(2-4): 159-65.
  33. Tigno XT, de Guzman F, Flora AM. Phytochemical analysis and hemodynamic actions of *Artemisia vulgaris* L. *Clin Hemorheol Microcirc* 2000; 23(2-4): 167-75.
  34. Singh R, De S, Belkheir A. *Avena sativa* (Oat), a potential nutraceutical and therapeutic agent: An overview. *Crit Rev Food Sci Nutr* 2013; 53(2): 126-44.
  35. Gasparis S, Nadolska-Orczyk A. Oat (*Avena sativa* L.). *Methods Mol Biol* 2015; 1223: 143-53.
  36. Criquet M, Roure R, Dayan L, Nollent V, Bertin C. Safety and efficacy of personal care products containing colloidal oatmeal. *Clin Cosmet Investig Dermatol* 2012; 5: 183-93.
  37. Chan E. Displacement of bilirubin from albumin by berberine. *Biol Neonate* 1993; 63(4): 201-8.
  38. Khalid Nusier M, Nayef Bataineh H, Mohydeen Bataineh Z, Daradka HM. Effect of frankincense (*Boswellia thurifera*) on reproductive system in adult male rat. *J Health Sci* 2007; 53(4): 365-70.
  39. Kulkarni RR, Patki PS, Jog VP, Gandage SG, Patwardhan B. Treatment of osteoarthritis with a herbomineral formulation: A double-blind, placebo-controlled, cross-over study. *J Ethnopharmacol* 1991; 33(1-2): 91-5.
  40. Jellin JF, Gregory P, Batz F. *Natural Medicines Comprehensive Database*. Stockton, CA: Therapeutic Research Faculty; 2000. p. 1521.
  41. Shipochliev T. Uterotonic action of extracts from a group of medicinal plants. *Vet Med Nauki* 1981; 18(4): 94-8.
  42. Mennitti LV, Oyama LM, de Oliveira JL, Hachul AC, Santamarina AB, de Santana AA, et al. Oligofructose supplementation during pregnancy and lactation impairs offspring development and



- alters the intestinal properties of 21-d-old pups. *Lipids Health Dis* 2014; 13: 26.
43. Kulkarni TR, Kothekar MA, Mateenuddin M. Study of anti-fertility effect of lemon seeds (*Citrus limonum*) in female albino mice. *Indian J Physiol Pharmacol* 2005; 49(3): 305-12.
  44. Nwoha PU. The immobilization of all spermatozoa in vitro by bitter lemon drink and the effect of alkaline pH. *Contraception* 1992; 46(6): 537-42.
  45. Yavari KP, Safajou F, Shahnazi M, Nazemiyeh H. The effect of lemon inhalation aromatherapy on nausea and vomiting of pregnancy: A double-blinded, randomized, controlled clinical trial. *Iran Red Crescent Med J* 2014; 16(3): e14360.
  46. Kuhn MA, Winston D. *Winston & Kuhn's herbal therapy and supplements: A scientific and traditional approach*. 2<sup>nd</sup> ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2012.
  47. Shah AH, Al-Shareef AH, Ageel AM, Qureshi S. Toxicity studies in mice of common spices, *Cinnamomum zeylanicum* bark and *Piper longum* fruits. *Plant Foods Hum Nutr* 1998; 52(3): 231-9.
  48. Akour A, Kasabri V, Afifi FU, Bulatova N. The use of medicinal herbs in gynecological and pregnancy-related disorders by Jordanian women: A review of folkloric practice vs. evidence-based pharmacology. *Pharm Biol* 2016; 54(9): 1901-18.
  49. Lindhout D, Omtzigt JG. Pregnancy and the risk of teratogenicity. *Epilepsia* 1992; 33(Suppl 4): S41-S48.
  50. Cameron EL. Pregnancy and olfaction: a review. *Front Psychol* 2014; 5: 67.
  51. Lepik K. Safety of herbal medications in pregnancy. *Can Pharm J* 1997; 130(3): 29-33.
  52. Dugoua JJ, Seely D, Perri D, Cooley K, Forelli T, Mills E, et al. From type 2 diabetes to antioxidant activity: A systematic review of the safety and efficacy of common and cassia cinnamon bark. *Can J Physiol Pharmacol* 2007; 85(9): 837-47.
  53. Zaidi SF, Aziz M, Muhammad JS, Kadowaki M. Review: Diverse pharmacological properties of *Cinnamomum cassia*: A review. *Pak J Pharm Sci* 2015; 28(4): 1433-8.
  54. Hajhoseini L. Importance of optimal fiber consumption during pregnancy. *International Journal of Women's Health and Reproduction Sciences* 2013; 1(3): 76-9.
  55. Arguelles N, Alvarez-Gonzalez I, Chamorro G, Madrigal-Bujaidar E. Protective effect of grapefruit juice on the teratogenic and genotoxic damage induced by cadmium in mice. *J Med Food* 2012; 15(10): 887-93.
  56. Rasch V. Cigarette, alcohol, and caffeine consumption: Risk factors for spontaneous abortion. *Acta Obstet Gynecol Scand* 2003; 82(2): 182-8.
  57. Wisborg K, Kesmodel U, Bech BH, Hedegaard M, Henriksen TB. Maternal consumption of coffee during pregnancy and stillbirth and infant death in first year of life: prospective study. *BMJ* 2003; 326(7386): 420.
  58. Evreklioglu C, Sari I, Alasehirli B, Guldur E, Cengiz B, Balat Z, et al. High dose of caffeine administered to pregnant rats causes histopathological changes in the cornea of newborn pups. *Med Sci Monit* 2003; 9(5): BR168-BR173.
  59. Ajarem JS, Ahmad M. Teratopharmacological and behavioral effects of coffee in mice. *Acta Physiol Pharmacol Bulg* 1996; 22(2): 51-61.
  60. Palm PE, Arnold EP, Nick MS, Valentine JR, Doerfler TE. Two-year toxicity/carcinogenicity study of fresh-brewed coffee in rats initially exposed in utero. *Toxicol Appl Pharmacol* 1984; 74(3): 364-82.
  61. Munoz L, Keen CL, Lonnerdal B, Dewey KG. Coffee intake during pregnancy and lactation in rats: Maternal and pup hematological parameters and liver iron, zinc and copper concentration. *J Nutr* 1986; 116(7): 1326-33.
  62. Briggs GG, Freeman RK, Yaffe SJ. *Drugs in Pregnancy and Lactation: A Reference Guide to Fetal and Neonatal Risk*. Philadelphia, PA: Lippincott Williams & Wilkins; 2011.
  63. Ferrazzano GF, Amato I, Ingenito A, De Natale A, Pollio A. Anti-cariogenic effects of polyphenols from plant stimulant beverages (cocoa, coffee, tea). *Fitoterapia* 2009; 80(5): 255-62.
  64. Brinker FJ. *The toxicology of botanical medicines*. Sandy, OR: Eclectic Medical Publications; 2000.
  65. Dwyer PL, O'Reilly M. Recurrent urinary tract infection in the female. *Curr Opin Obstet Gynecol* 2002; 14(5): 537-43.
  66. Nordeng H, Havnen GC. Use of herbal drugs in pregnancy: a survey among 400 Norwegian women. *Pharmacoepidemiol Drug Saf* 2004; 13(6): 371-80.
  67. Foster S. *Tyler's honest herbal: A sensible guide to the use of herbs and related remedies*. London, UK: Routledge; 2012.
  68. Newall CA, Anderson LA, Phillipson JD. *Herbal medicines: A guide for health-care professionals*. London, UK: Pharmaceutical Press; 1996.
  69. Mills S, Bone K. *Principles and practice of phytotherapy: Modern herbal medicine*. London, UK: Churchill Livingstone; 2000.
  70. Yu ZY, Guo L, Wang B, Kang LP, Zhao ZH, Shan YJ, et al. Structural requirement of spirostanol glycosides for rat uterine contractility and mode of their synergism. *J Pharm Pharmacol* 2010; 62(4): 521-9.
  71. Seely D, Dugoua JJ, Perri D, Mills E, Koren G. Safety and efficacy of panax ginseng during pregnancy and lactation. *Can J Clin Pharmacol* 2008; 15(1): e87-e94.

72. Chan LY, Chiu PY, Lau TK. Embryotoxicity study of ginsenoside Rc and Rein in vitro rat whole embryo culture. *Reprod Toxicol* 2004; 19(1): 131-4.
73. Choi J, Kim TH, Choi TY, Lee MS. Ginseng for health care: A systematic review of randomized controlled trials in Korean literature. *PLoS One* 2013; 8(4): e59978.
74. Alsaad AM, Fox C, Koren G. Toxicology and teratology of the active ingredients of professional therapy MuscleCare products during pregnancy and lactation: A systematic review. *BMC Complement Altern Med* 2015; 15: 40.
75. Dreisinger N, Zane D, Etwaru K. A poisoning of topical importance. *Pediatr Emerg Care* 2006; 22(12): 827-9.
76. Darben T, Cominos B, Lee CT. Topical eucalyptus oil poisoning. *Australas J Dermatol* 1998; 39(4): 265-7.
77. Duke JA. Handbook of medicinal herbs. Boca Raton, FL: CRC Press; 2002.
78. Zehra U, Tahir M, Lone KP. Ginkgo biloba induced malformations in mice. *J Coll Physicians Surg Pak* 2010; 20(2): 117-21.
79. Dugoua JJ, Mills E, Perri D, Koren G. Safety and efficacy of ginkgo (*Ginkgo biloba*) during pregnancy and lactation. *Can J Clin Pharmacol* 2006; 13(3): e277-e284.
80. Leung AY. Encyclopedia of common natural ingredients used in food, drugs, and cosmetics. New York, NY: Wiley; 1980.
81. Natarajan SS, Xu C, Bae H, Caperna TJ, Garrett WM. Characterization of storage proteins in wild (*Glycine soja*) and cultivated (*Glycine max*) soybean seeds using proteomic analysis. *J Agric Food Chem* 2006; 54(8): 3114-20.
82. Benaiges A, Marcet P, Armengol R, Betes C, Girones E. Study of the refirming effect of a plant complex. *Int J Cosmet Sci* 1998; 20(4): 223-33.
83. Strandberg TE, Jarvenpaa AL, Vanhanen H, McKeigue PM. Birth outcome in relation to licorice consumption during pregnancy. *Am J Epidemiol* 2001; 153(11): 1085-8.
84. Rees WD, Rhodes J, Wright JE, Stamford LF, Bennett A. Effect of deglycyrrhized licorice on gastric mucosal damage by aspirin. *Scand J Gastroenterol* 1979; 14(5): 605-7.
85. Iyare EE, Iyare FE. Maternal consumption of aqueous extract of hibiscus sabdariffa during pregnancy attenuates pregnancy weight gain and postpartum weight loss. *Afr J Biomed Res* 2007; 10(3): 257-61.
86. Iyare EE, Adegoke OA. Maternal consumption of an aqueous extract of *Hibiscus sabdariffa* during lactation accelerates postnatal weight and delays onset of puberty in female offspring. *Niger J Physiol Sci* 2008; 23(1-2): 89-94.
87. Henley DV, Lipson N, Korach KS, Bloch CA. Prepubertal gynecomastia linked to lavender and tea tree oils. *N Engl J Med* 2007; 356(5): 479-85.
88. Basch E, Foppa I, Liebowitz R, Nelson J, Smith M, Sollars D, et al. Lavender (*Lavandula angustifolia* Miller). *J Herb Pharmacother* 2004; 4(2): 63-78.
89. Ernst E. Herbal medicinal products during pregnancy: Are they safe? *BJOG* 2002; 109(3): 227-35.
90. Farnsworth NR, Bingel AS, Cordell GA, Crane FA, Fong HS. Potential value of plants as sources of new antifertility agents II. *J Pharm Sci* 1975; 64(5): 717-54.
91. Casanova M, You L, Gaido KW, Archibeque-Engle S, Janszen DB, Heck HA. Developmental effects of dietary phytoestrogens in Sprague-Dawley rats and interactions of genistein and daidzein with rat estrogen receptors alpha and beta in vitro. *Toxicol Sci* 1999; 51(2): 236-44.
92. Aqel M, Shaheen R. Effects of the volatile oil of *Nigella sativa* seeds on the uterine smooth muscle of rat and guinea pig. *J Ethnopharmacol* 1996; 52(1): 23-6.
93. Salarinia R, Rakhshandeh H, Oliaee D, Gul GS, Ghorbani A. Safety evaluation of Phytovagex, a pessary formulation of *Nigella sativa*, on pregnant rats. *Avicenna J Phytomed* 2016; 6(1): 117-23.
94. Kenny FS, Pinder SE, Ellis IO, Gee JM, Nicholson RI, Bryce RP, et al. Gamma linolenic acid with tamoxifen as primary therapy in breast cancer. *Int J Cancer* 2000; 85(5): 643-8.
95. Dove D, Johnson P. Oral evening primrose oil: Its effect on length of pregnancy and selected intrapartum outcomes in low-risk nulliparous women. *J Nurse Midwifery* 1999; 44(3): 320-4.
96. McFarlin BL, Gibson MH, O'Rear J, Harman P. A national survey of herbal preparation use by nurse-midwives for labor stimulation. Review of the literature and recommendations for practice. *J Nurse Midwifery* 1999; 44(3): 205-16.
97. Brinker FJ. Herb contraindications and drug interactions: With appendices addressing specific conditions and medicines. Sandy, OR: Eclectic Medical Publications; 1998.
98. Berdai MA, Labib S, Harandou M. Peganum harmala L. Intoxication in a pregnant woman. *Case Rep Emerg Med* 2014; 2014: 783236.
99. Blotman F, Maheu E, Wulwik A, Caspard H, Lopez A. Efficacy and safety of avocado/soybean unsaponifiables in the treatment of symptomatic osteoarthritis of the knee and hip. A prospective, multicenter, three-month, randomized, double-blind, placebo-controlled trial. *Rev Rhum Engl Ed* 1997; 64(12): 825-34.
100. Rezazad M, Farokhi F. Protective effect of *Petroselinum crispum* extract in abortion using

- prostadin-induced renal dysfunction in female rats. *Avicenna J Phytomed* 2014; 4(5): 312-9.
101. Greenhalf JO, Leonard HS. Laxatives in the treatment of constipation in pregnant and breast-feeding mothers. *Practitioner* 1973; 210(256): 259-63.
  102. Ghahramani L, Hosseini SV, Rahimikazerooni S, Bananzadeh AM, Jahromi B, Samsam A. The effect of oral psyllium herbal laxative powder in prevention of hemorrhoids and anal fissure during pregnancy, a randomized double blind clinical trial. *Ann Colorectal Res* 2013; 1(1): 23-7.
  103. McRorie JW, Daggy BP, Morel JG, Diersing PS, Miner PB, Robinson M. Psyllium is superior to docusate sodium for treatment of chronic constipation. *Aliment Pharmacol Ther* 1998; 12(5): 491-7.
  104. Cheskin LJ, Kamal N, Crowell MD, Schuster MM, Whitehead WE. Mechanisms of constipation in older persons and effects of fiber compared with placebo. *J Am Geriatr Soc* 1995; 43(6): 666-9.
  105. Londonkar R, Nayaka H. Evaluation of anti implantation and abortifacient properties of portulaca oleracea l. In albino rats. *Int J Pharma Bio Sci* 2011; 2(4): 501-8.
  106. Hooman N, Mojab F, Nickavar B, Pouryousefi-Kermani P. Diuretic effect of powdered *Cerasus avium* (cherry) tails on healthy volunteers. *Pak J Pharm Sci* 2009; 22(4): 381-3.
  107. West T, Atzeva M, Holtzman DM. Pomegranate polyphenols and resveratrol protect the neonatal brain against hypoxic-ischemic injury. *Dev Neurosci* 2007; 29(4-5): 363-72.
  108. Shidfar F, Rahideh ST, Rajab A, Khandozi N, Hosseini S, Shidfar S, et al. The effect of sumac (*Rhus coriaria* L.) Powder on Serum Glycemic Status, ApoB, ApoA-I and total antioxidant capacity in type 2 diabetic patients. *Iran J Pharm Res* 2014; 13(4): 1249-55.
  109. Chaube S, Swinyard CA. Teratological and toxicological studies of alkaloidal and phenolic compounds from *Solanum tuberosum* L. *Toxicol Appl Pharmacol* 1976; 36(2): 227-37.
  110. Klebanoff MA, Levine RJ, DerSimonian R, Clemens JD, Wilkins DG. Maternal serum paraxanthine, a caffeine metabolite, and the risk of spontaneous abortion. *N Engl J Med* 1999; 341(22): 1639-44.
  111. Sindos M, Pisal N, Michala S. Consumption of coffee during pregnancy: Authors should adjust for history of drug abuse. *BMJ* 2003; 326(7401): 1268.
  112. Balat O, Balat A, Ugur MG, Pence S. The effect of smoking and caffeine on the fetus and placenta in pregnancy. *Clin Exp Obstet Gynecol* 2003; 30(1): 57-9.
  113. Bracken MB, Triche EW, Belanger K, Hellenbrand K, Leaderer BP. Association of maternal caffeine consumption with decrements in fetal growth. *Am J Epidemiol* 2003; 157(5): 456-66.
  114. Yazdy MM, Tinker SC, Mitchell AA, Demmer LA, Werler MM. Maternal tea consumption during early pregnancy and the risk of spina bifida. *Birth Defects Res A Clin Mol Teratol* 2012; 94(10): 756-61.
  115. Yarnell E, Abascal K. Dandelion (*Taraxacum officinale* and *T. mongolicum*). *Integrative Medicine* 2009; 8(2): 35-8.
  116. Sharma PV. *Classical Uses of Medicinal Plants*. Varanasi, India: Chaukhambha Visvabharati; 1996.
  117. Youdim KA, Deans SG. Effect of thyme oil and thymol dietary supplementation on the antioxidant status and fatty acid composition of the ageing rat brain. *Br J Nutr* 2000; 83(1): 87-93.
  118. Blumenthal M. *Herbal medicine: Expanded commission e monographs*. Newton, MA: Integrative Medicine Communications; 2000.
  119. Heinonen OP, Slone D, Shapiro S. *Birth defects and drugs in pregnancy*. Littleton, MA: Publishing Sciences Group; 1977.
  120. Korman SH, Cohen E, Preminger A. Pseudo-maple syrup urine disease due to maternal prenatal ingestion of fenugreek. *J Paediatr Child Health* 2001; 37(4): 403-4.
  121. Abdo MS, al-Kafawi AA. Experimental studies on the effect of *Trigonella foenum-graecum*. *Planta Med* 1969; 17(1): 14-8.
  122. Edirne T, Arica SG, Gucuk S, Yildizhan R, Kolusari A, Adali E, et al. Use of complementary and alternative medicines by a sample of Turkish women for infertility enhancement: A descriptive study. *BMC Complement Altern Med* 2010; 10: 11.
  123. Jalili C, Salahshoor MR, Naseri A. Protective effect of *Urtica dioica* L against nicotine-induced damage on sperm parameters, testosterone and testis tissue in mice. *Iran J Reprod Med* 2014; 12(6): 401-8.
  124. Bercovich E, Saccomanni M. Analysis of the results obtained with a new phytotherapeutic association for LUTS versus control. [corrected]. *Urologia* 2010; 77(3): 180-6.
  125. Azarnia M, Ejtemaee-Mehr S, Shakoob A. Effects of vitex agnus castus on mice fetus development. *Acta Med Iran* 2007; 45(4): 263-70.
  126. Rani A, Sharma A. The genus *Vitex*: A review. *Pharmacogn Rev* 2013; 7(14): 188-98.
  127. Ladeji O, Udoh FV, Okoye ZS. Activity of aqueous extract of the bark of *Vitex doniana* on uterine muscle response to drugs. *Phyther Res* 2005; 19(9): 804-6.
  128. Daniele C, Thompson Coon J, Pittler MH, Ernst E. *Vitex agnus castus*: A systematic review of adverse events. *Drug Saf* 2005; 28(4): 319-32.
  129. Takahashi T, Yokoo Y, Inoue T, Ishii A. Toxicological studies on procyanidin B-2 for external application as a hair growing agent. *Food Chem Toxicol* 1999; 37(5): 545-52.

130. Erexson GL. Lack of in vivo clastogenic activity of grape seed and grape skin extracts in a mouse micronucleus assay. *Food Chem Toxicol* 2003; 41(3): 347-50.
131. Ahmad MK, Mahdi AA, Shukla KK, Islam N, Rajender S, Madhukar D, et al. *Withania somnifera* improves semen quality by regulating reproductive hormone levels and oxidative stress in seminal plasma of infertile males. *Fertil Steril* 2010; 94(3): 989-96.
132. Malviya N, Jain S, Gupta VB, Vyas S. Recent studies on aphrodisiac herbs for the management of male sexual dysfunction--a review. *Acta Pol Pharm* 2011; 68(1): 3-8.
133. Fischer-Rasmussen W, Kjaer SK, Dahl C, Asping U. Ginger treatment of hyperemesis gravidarum. *Eur J Obstet Gynecol Reprod Biol* 1991; 38(1): 19-24.
134. Keating A, Chez RA. Ginger syrup as an antiemetic in early pregnancy. *Altern Ther Health Med* 2002; 8(5): 89-91.
135. Portnoi G, Chng LA, Karimi-Tabesh L, Koren G, Tan MP, Einarson A. Prospective comparative study of the safety and effectiveness of ginger for the treatment of nausea and vomiting in pregnancy. *Am J Obstet Gynecol* 2003; 189(5): 1374-7.
136. Weidner MS, Sigwart K. Investigation of the teratogenic potential of a zingiber officinale extract in the rat. *Reprod Toxicol* 2001; 15(1): 75-80.
137. Nakamura H, Yamamoto T. Mutagen and anti-mutagen in ginger, *Zingiber officinale*. *Mutat Res* 1982; 103(2): 119-26.
138. Nakamura H, Yamamoto T. The active part of the [6]-gingerol molecule in mutagenesis. *Mutat Res* 1983; 122(2): 87-94.
139. Wilkinson JM. Effect of ginger tea on the fetal development of Sprague-Dawley rats. *Reprod Toxicol* 2000; 14(6): 507-12.
140. Jewell D, Young G. Interventions for nausea and vomiting in early pregnancy. *Cochrane Database Syst Rev* 2003; (4): CD000145.
141. Viljoen E, Visser J, Koen N, Musekiwa A. A systematic review and meta-analysis of the effect and safety of ginger in the treatment of pregnancy-associated nausea and vomiting. *Nutr J* 2014; 13: 20.
142. Gupta RB, Sharma S, Sharma J, Goyal R. Study on the physicochemical characters of fruits of some wild and cultivated forms/spp." *Zizpi hus spp. Haryana J Horticult Sci* 2004; 33(3/4): 167-69.
143. Gupta M, Mazumder UK, Vamsi ML, Sivakumar T, Kandar CC. Anti-steroidogenic activity of the two Indian medicinal plants in mice. *J Ethnopharmacol* 2004; 90(1): 21-5.
144. Wiesbaden Wagner H, Wiesnauer M. *Phytotherapie: Phytopharmaka und pflanzliche Homöopathika*. Jena, Germany: Gustav Fischer Verlag; 1995.
145. Isaacsohn JL, Moser M, Stein EA, Dudley K, Davey JA, Liskov E, et al. Garlic powder and plasma lipids and lipoproteins: A multicenter, randomized, placebo-controlled trial. *Arch Intern Med* 1998; 158(11): 1189-94.
146. Holzgartner H, Schmidt U, Kuhn U. Comparison of the efficacy and tolerance of a garlic preparation vs. bezafibrate. *Arzneimittelforschung* 1992; 42(12): 1473-7.
147. Fleming T. *PDR for Herbal Medicine*. 1st ed. Montvale, NJ: Medical Economics Co; 1998.
148. Calomme M, Pieters L, Vlietinck A, Vanden Berghe D. Inhibition of bacterial mutagenesis by Citrus flavonoids. *Planta Med* 1996; 62(3): 222-6.
149. Stapleton AE. Cranberry-containing products are associated with a protective effect against urinary tract infections. *Evid Based Med* 2013; 18(3): 110-1.
150. Oh HG, Lee HY, Seo MY, Kang YR, Kim JH, Park JW, et al. Effects of *Ficus carica* paste on constipation induced by a high-protein feed and movement restriction in beagles. *Lab Anim Res* 2011; 27(4): 275-81.
151. Ravn-Haren G, Dragsted LO, Buch-Andersen T, Jensen EN, Jensen RI, Nemeth-Balogh M, et al. Intake of whole apples or clear apple juice has contrasting effects on plasma lipids in healthy volunteers. *Eur J Nutr* 2013; 52(8): 1875-89.
152. Anderson JW, Allgood LD, Turner J, Oeltgen PR, Daggly BP. Effects of psyllium on glucose and serum lipid responses in men with type 2 diabetes and hypercholesterolemia. *Am J Clin Nutr* 1999; 70(4): 466-73.
153. Zhang Y, Neogi T, Chen C, Chaisson C, Hunter DJ, Choi HK. Cherry consumption and decreased risk of recurrent gout attacks. *Arthritis Rheum* 2012; 64(12): 4004-11.
154. Blaschek W, Hänsel R, Keller K, Reichling J., Rimpler H, Schneider G. *Hagers Handbuch der Pharmazeutischen Praxis*. Berlin, Germany: Springer-Verlag Berlin Heidelberg; 1990.
155. Kalume DE, Sousa MV, Morhy L. Purification, characterization, sequence determination, and mass spectrometric analysis of a trypsin inhibitor from seeds of the Brazilian tree *Dipteryx alata* (Leguminosae). *J Protein Chem* 1995; 14(8): 685-93.
156. Henriët JP. Venous-lymphatic insufficiency. 4,729 patients undergoing hormonal and procyanidol oligomer therapy. *Phlebologie* 1993; 46(2): 313-25.
157. Tebib K, Rouanet JM, Besancon P. Effect of grape seed tannins on the activity of some rat intestinal enzyme activities. *Enzyme Protein* 1994; 48(1): 51-60.
158. Muller JL, Clauson KA. Pharmaceutical considerations of common herbal medicine. *Am J Managed Care* 1997; 3(11): 1753-70.
159. Rumbold A, Duley L, Crowther CA, Haslam RR. Antioxidants for preventing pre-eclampsia. *Cochrane Database Syst Rev* 2008; (1): CD004227.
160. Mistry HD, Williams PJ. The importance of

- antioxidant micronutrients in pregnancy. *Oxid Med Cell Longev* 2011; 2011.
161. Jenkins C, Wilson R, Roberts J, Miller H, McKillop JH, Walker JJ. Antioxidants: Their role in pregnancy and miscarriage. *Antioxid Redox Signal* 2000; 2(3): 623-8.
162. Park H. Effects of antioxidants and oxidative stress on pregnancy and infant growth: Korean perspectives. In: Preedy VR, Editor. *Handbook of growth and growth monitoring in health and disease*. Berlin, Germany: Springer Science & Business Media; 2011. p. 1585-98.
163. Rouhi-Boroujeni H, Heidarian E, Rouhi-

Boroujeni H, Deris F, Rafieian-Kopaei M. Medicinal plants with multiple effects on cardiovascular diseases: A systematic review. *Curr Pharm Des* 2017; 23(7): 999-1015.

**How to cite this article:** Rouhi-Boroujeni H, Heidarian E, Rouhi-Boroujeni H, Khoddami M, Gharipour M, Rafieian-Kopaei M. **Use of lipid-lowering medicinal herbs during pregnancy: A systematic review on safety and dosage.** *ARYA Atheroscler* 2017; 13(3): 135-55.